



September 5, 2019

Public Health Advisory Severe Pulmonary Disease Associated with Using E-Cigarette Products

The City of Amarillo Public Health Department is issuing this health advisory to provide recommendations for clinicians and the public about severe pulmonary disease associated with using e-cigarette products. The following is an overview of the 8/30/19 Health Advisory issued by the Centers for Disease Control and Prevention (CDC). The original document can be found at the following link: <https://emergency.cdc.gov/han/han00421.asp>.

General Background

E-cigarettes are known by many different names and come in a variety of shapes, sizes and device types. Devices may resemble cigarettes, USB flash drives, pens, flashlights or have unique shapes. Use of e-cigarettes is sometimes referred to as “vaping” or “juuling.”

E-cigarettes can contain harmful or potentially harmful substances, including nicotine, heavy metals, volatile organic compounds, and cancer-causing chemicals. Additionally, some e-cigarette products are used to deliver illicit substances; may be acquired from unknown or unauthorized sources; and may be modified for uses that could increase their potential for harm to the user.

Outbreak background

As of August 27, 2019, 215 possible cases have been reported from 25 states and additional reports of pulmonary illness are under investigation. One patient (in Illinois) with a history of recent e-cigarette use was hospitalized with severe pulmonary disease and subsequently died. Although the etiology of e-cigarette-associated pulmonary disease is undetermined, epidemiologic investigations in affected states are ongoing to better characterize the exposures, demographic, clinical, and laboratory features and behaviors of patients. All patients have reported using e-cigarette products. The exact number is currently unknown, but many patients have reported using e-cigarettes containing cannabinoid products such as THC or CBD.

Based on reports from several states, patients have experienced respiratory symptoms (cough, shortness of breath, or chest pain), and some have also experienced gastrointestinal symptoms (nausea, vomiting, or diarrhea) or non-specific constitutional symptoms (fatigue, fever, or weight loss). Symptoms typically develop over a period of days but sometimes can manifest over several weeks. Gastrointestinal symptoms sometimes preceded respiratory symptoms. Fever, tachycardia, and elevated white blood cell count have been reported in the absence of an identifiable infectious disease. Many patients have sought initial care in ambulatory settings, some with several visits, before hospital admission.

Radiologic findings have varied and are not present in all patients upon initial presentation. Bilateral pulmonary infiltrates and diffuse ground-glass opacities have been reported. Many patients required supplemental oxygen, some required assisted ventilation and oxygenation, and some were intubated. Some patients have been treated with corticosteroids with demonstrated improvement. Antimicrobial therapy alone has not consistently been associated with clinical improvement. Assessment for infectious

City of Amarillo Department of Public Health

Amarillo Bi-City, Bi-County Health District
1000 Martin Road, Amarillo, TX 79107
(806) 378-6300
amarillopublichealth.org



etiologies has been completed in many patients without an identified infectious cause. Several patients from one state have been diagnosed with lipid pneumonia based on clinical presentation and detection of lipids within bronchoalveolar lavage samples stained specifically to detect oil.

All patients have reported using e-cigarette products and the symptom onset has ranged from a few days to several weeks after e-cigarette use. Within two states, recent inhalation of cannabinoid products, THC or cannabidiol, have been reported in many of the patients. To date, no single substance or e-cigarette product has been consistently associated with illness. CDC is working closely with state health departments to facilitate collecting product specimens for testing at the U.S. FDA Forensic Chemistry Center.

Recommendations for Clinicians

1. Report cases of severe pulmonary disease of unclear etiology and a history of e-cigarette product use within the past 90 days to the City of Amarillo Department of Public Health – Laurie Burton, Communicable Disease Program Manager at 806-378-6321 or Kirstin Williams, Epidemiologist at 806-378-6353. Reporting of cases may help CDC and state health departments determine the cause or causes of these pulmonary illnesses.
2. Ask all patients who report e-cigarette product use within the last 90 days about signs and symptoms of pulmonary illness.
3. If e-cigarette product use is suspected as a possible etiology of a patient’s severe pulmonary disease, obtain detailed history regarding:
 - Substance(s) used: nicotine, cannabinoids (e.g., marijuana, THC, THC concentrates, CBD, CBD oil, synthetic cannabinoids [e.g., K2 or spice], hash oil, Dank vapes), flavors, or other substances
 - Substance source(s): commercially available liquids (i.e., bottles, cartridges, or pods), homemade liquids, and re-use of old cartridges or pods with homemade or commercially bought liquids
 - Device(s) used: manufacturer; brand name; product name; model; serial number of the product, device, or e-liquid; if the device can be customized by the user; and any product modifications by the user (e.g., exposure of the atomizer or heating coil)
 - Where the product(s) were purchased
 - Method of substance use: aerosolization, dabbing, or dripping
 - Other potential cases: sharing e-cigarette products (devices, liquids, refill pods, or cartridges) with others
4. Determine if any remaining product, including devices and liquids, are available for testing. Testing can be coordinated with the local or state health departments.
5. Consider all possible causes of illness in patients reporting respiratory and gastrointestinal symptoms and of e-cigarette product use. Evaluate and treat for other possible causes of illness (e.g., infectious, rheumatologic, neoplastic) as clinically indicated. Consider consultation with specialists (pulmonary, infectious disease, critical care, medical toxicology) as appropriate.
6. Clinical improvement of patients with severe pulmonary disease associated with e-cigarette use has been reported with the use of corticosteroids. The decision to use corticosteroids should be made on a case-by-case basis based on risks and benefits and the likelihood of other etiologies.
7. Lipid pneumonia associated with inhalation of lipids in aerosols generated by e-cigarettes has been reported based on the detection of lipid-laden alveolar macrophages obtained by bronchoalveolar

City of Amarillo Department of Public Health

Amarillo Bi-City, Bi-County Health District
1000 Martin Road, Amarillo, TX 79107
(806) 378-6300
amarillopublichealth.org



- lavage (BAL) and lipid staining (e.g., oil red O). The decision about whether to perform a BAL should be based on individual clinical circumstances.
8. Lung biopsies have been performed on some patients. If a lung biopsy is obtained, lipid staining may be considered during pathologic examination, and is best performed on fresh tissue. Routine pathology tissue processing (including formalin-fixation and paraffin-embedding) can remove lipids. Conducting routine tissue processing and histopathologic evaluation is still important. Consider consultation with specialists in pulmonary medicine and pathology to help inform any evaluation plan.
 9. Patients who have received treatment for severe pulmonary disease related to e-cigarette product use should undergo follow-up evaluation as clinically indicated to monitor pulmonary function.

Recommendations for the Public

1. While this investigation is ongoing, if you are concerned about these specific health risks, consider refraining from using e-cigarette products.
2. Regardless of the ongoing investigation, anyone who uses e-cigarette products should not buy these products off the street (e.g., e-cigarette products with THC, other cannabinoids) and should not modify e-cigarette products or add any substances to these products that are not intended by the manufacturer.
3. Regardless of the ongoing investigation, e-cigarette products should not be used by youth, young adults, pregnant women, as well as adults who do not currently use tobacco products. If you use e-cigarette products, monitor yourself for symptoms (e.g., cough, shortness of breath, chest pain) and promptly seek medical attention if you have concerns about your health. CDC and FDA will continue to advise and alert the public as more information becomes available.
4. Adult smokers who are attempting to quit should use evidence-based treatments, including counseling and FDA-approved medications. If you who need help quitting tobacco products, including e-cigarettes, contact your doctor. You can also call the City of Amarillo Department of Public Health at 806-378-6335 for a schedule of Tobacco Cessation classes offered through Tobacco Free Amarillo.
5. If you are concerned about harmful effects from e-cigarette products, call your local poison control center at: 1-800-222-1222.
6. We encourage the public to submit detailed reports of any unexpected tobacco or e-cigarette-related health or product issues to the FDA via the online Safety Reporting Portal: <https://www.safetyreporting.hhs.gov/external/icon>.

City of Amarillo Department of Public Health

Amarillo Bi-City, Bi-County Health District
1000 Martin Road, Amarillo, TX 79107
(806) 378-6300
amarillopublichealth.org